

NEBRASKA

EMERGENCY MEDICAL SERVICES

MODEL PROTOCOLS

SERVICE NAME

With the approval of the Physician Medical Director, the service has adopted the following protocols

Date Approved

Date Reviewed

Physician Medical Director

Agency Head

TABLE OF CONTENTS

Subject	Page
Introduction	iii
General Operations	iv
General Principles	ix
Protocols	
<i>Adult Cardiovascular Emergencies</i>	
Cardiopulmonary Arrest	1
Adult Cardiac Arrest Algorithm V-fib	2
Adult Cardiac Arrest Algorithm PEA	3
Adult Cardiac Arrest Algorithm Asystole	4
AED	5
Post Cardiac Arrest – Return of Pulse	6
Discontinue CPR	7
General Cardiac Dysrhythmia	8
Advanced Cardiac Dysrhythmia	9
Chest Pain	11
Pulmonary Edema	12
Cardiogenic Shock	13
<i>Difficulty Breathing</i>	
Acute Allergic Reaction / Anaphylaxis	14
Asthma	16
COPD	17
Respiratory Infection	18
Adult Epiglottitis	19
<i>Adult Medical Emergencies</i>	
Adult Upper Airway Obstruction	20
Non-Traumatic Alter or Decreased Level of Consciousness	
Diabetes Mellitus –Hypoglycemia	21
Hyperglycemia	22
Altered Mental Status	23
Cerebrovascular Accident	24
Seizure Disorder	25
Hypothermia	26
Hyperthermia	27
Non-Traumatic Abdominal Pain	28
Upper and Lower GI Bleed	29
Hypotension in Absence of Trauma	30
<i>Obstetric and Gynecological Emergencies</i>	
Imminent Delivery	31
Newborn Care	32
Birth Complications	33
Hypertensive Disorders of Pregnancy	34

Vaginal Bleeding	35
<i>Acute Traumatic Emergencies</i>	
General Trauma Management	36
Trauma System	37
Head/Facial Injuries	39
Soft Tissue Neck Injuries	40
Chest Injuries	41
Abdominal/Pelvic Injuries	42
Extremity Injuries	43
Ingested Poisons	44
Specific Agents/Toxins	45 - 48
Toxic Inhalation	49
Burns	50
Snake Bite	51
<i>Pediatrics</i>	
Cardiopulmonary Arrest	52
Pediatric Cardiac Arrest Algorithm V-Fib	53
Pediatric Cardiac Arrest Algorithm PEA	54
Pediatric Cardiac Arrest Algorithm Asystole	55
Pediatric Post Cardiac Arrest – Return of Pulse	56
Pediatric General Cardiac Dysrhythmia	57
Pediatric Advanced Cardiac Dysrhythmia	58
Pediatric Difficulty Breathing	60
Asthma	60
Laryngotracheobronchitis (Croup), Epiglottitis	61
Acute Allergic Reaction / Anaphylaxis	62
Pediatric Upper Airway Obstruction	63
Pediatric Seizures	64
<i>Appendix</i>	
Bronchodilator Protocol	65
Pain Management Protocol	66
Hospital to Hospital Transfers Protocol	67
Anti-Emetic Protocol	68
Rapid Sequence Intubation Protocol	69

Nebraska Emergency Medical Services
Protocols

**First Responder, Basic, EMT Intermediate, Paramedic
Introduction**

Purpose: The purpose of these protocols is to assure safe and effective intervention during the out-of-hospital phase of patient care. In consideration of the unique resources, needs, population, and geography of EMS in Nebraska, individual medical directors may choose to enhance or omit portions of these protocols in accordance with current medical practice and standards. Medical directors are responsible to ensure the EMS personnel using these protocols have the training and skills required, and perform quality assurance activities to assure these protocols are used appropriately. It is the hope of the Nebraska Health and Human Services, Board of Emergency Medical Services, that these protocols will serve as a standard throughout Nebraska's system. Ongoing review and update of these protocols is necessary to keep pace with interventions known to be effective in out-of-hospital care.

Authority: Out-of-hospital (OOH) emergency care provider personnel may only deliver emergency medical care as a member of a licensed emergency medical service. The emergency medical service must have a physician medical director who is responsible for the practice of the OOH emergency care provider personnel. All treatments and procedures performed by each OOH emergency care provider must be authorized by their service's physician medical director.

Protocols shall be approved, signed, and dated by the individual service's physician medical director prior to implementation.

Directions for Use:

All emergency care providers should start at the top of the page and proceed as far down the page as your level of certification **and** training permits.

Always conduct a scene size-up and observe body substance isolation precautions and only then perform the patient assessment and obtain the necessary information on all patients.

You may need to use more than one treatment protocol for any single patient, (known diabetic who has been in a motor vehicle crash).

The OOH emergency care provider must assess each patient and apply the correct treatment based on that assessment. All the treatment/interventions may not be required simply because the patient presents with a particular problem. Treatment must be applied based on the patient's condition and the provider's assessment. The provider is encouraged to contact medical control for advice if question about treatment arise.

GENERAL OPERATIONS

I. Scene Size-Up:

As you approach the scene, assure safety for yourself, your fellow responders and the patient. Establish and follow an Incident Command Plan.

II. BSI (Body Substance Isolation):

Prior to patient assessment, it is protocol policy to practice body substance isolation when caring for ALL patients. This includes washing hands after each patient care incident. Hands shall be washed even if gloves were worn or waterless soap was used. Note: This is comparable to Universal Precautions in a hospital setting.

This policy also applies to immediate disposal of needles and sharps in disposable, impervious containers. The practice of not recapping needles is highly encouraged.

III. Trauma Patients:

Once a trauma patient has been identified, follow the trauma system decision protocol for the identification of time critical injuries, the method of transport and the trauma facility resources necessary for treatment of those injuries.

IV. Use of Restraints

A. Indications:

A patient who needs to be transported for medical care, who appears to be in imminent danger to himself.

B. Protocol:

1. Check restraints as soon as applied and every 10 minutes thereafter to ensure no injury to extremities.
2. Once restrained, the patient is never to be left alone.
3. Written and verbal reports must completely document the necessity for the use of physical restraints.
4. Record condition of limbs before applying restraints and recheck and record condition on arrival at hospital.
5. Prevent asphyxia
 - a. Do not restrain patient prone (face down)
 - b. Do not restrain patient sandwiched between backboards, scoop stretchers, or other immobilization devices
 - c. Check and monitor any straps across the chest. Straps should secure the patient without restricting chest expansion.
 - d. Patient will not be "Hog Tied" (hands restrained behind back, feet restrained together and the two restrained attached together)

V. Transport Codes:

Code 1 (Category Green) – Minimal or no apparent disease or injury. Patient transported for examination

Code 2 (Category Yellow) – Obvious illness or injury, not serious but needs medical attention

Code 3 (Category Red) – Apparent serious injury or illness needing immediate medical attention

Code 99 – CPR in progress

Code 4 (Category Black) – Dead patient

TCC (Trauma Center Candidate) – Should be used in conjunction with code 3, and means the patient may be diverted directly to the trauma center that is appropriate for that area or region.

VI. Physician on Scene

When a physician is present on the scene and desires to direct the run, the EMT should:

- A. Inform the physician that if the physician directs the run, the physician must accompany the patient to the hospital. This must be documented on the patient care report.
- B. Inform the physician at the onset of the run that Out-of-Hospital personnel have strict legal guidelines and established protocols and they may not exceed those guidelines or protocols.
- C. Inform the physician that any procedure outside the legal guidelines for that level of care must be carried out by the physician.
- D. Out-of-hospital personnel have the right and obligation at any time there is gross deviation from the accepted protocol to contact the receiving hospital for further instruction. The physician on the scene should be informed that contact with the hospital is being made. If possible, it may be advisable to have the receiving hospital physician speak directly to the physician at the scene.

VII. “Do Not Resuscitate” (DNR) Orders or Requests and identification

A DNR is a written order by a physician that a patient should not be resuscitated or have CPR performed. A DNR must be signed by a physician, dated, and have the patient’s name. An out-of-hospital emergency care provider can honor a DNR. The out-of-hospital emergency care provider must be identified in the patient care report.

Requests for Do Not Resuscitate or perform CPR:

An OOH emergency care provider can honor an effective Living Will or Health Care Power of Attorney. This applies only to adults. OOH emergency care providers can presume the validity of either of these documents if signed in Nebraska. Documents from other states in compliance with that state’s laws are also valid in Nebraska.

Observation of an original or a photocopy of a living will or health care power of attorney must be documented in the patient care report. An OOH emergency care provider **shall not** honor a living will if there is no information or evidence that a physician has determined the patient is in a terminal condition or in a persistent vegetative state. If there is information or evidence that a physician has determined the patient is in a terminal condition or in a

persistent vegetative state, this information should be documented in the patient care report. The patient care report must contain information that the patient is an adult (is 19 or older or has been married).

An OOH emergency care provider can refuse to honor an effective Living Will or Health Care Power of Attorney decision if the decision is contrary to a formally adopted policy of the provider that is based on religious beliefs or sincerely held ethical or moral convictions. If a service providing out-of-hospital emergency care has such a policy approved by the Physician Medical Director, individual providers with such religious beliefs or ethical or moral convictions employed by or volunteering for that service may refuse to honor an effective Living Will or Health Care Power of Attorney. To the extent reasonably possible, the community in which this organization provides out-of-hospital emergency care shall be informed of the organization's formal policy. The appropriate person holding the Living Will or Health Care Power of Attorney, at the scene of the emergency, shall also be informed of the policy.

Discontinuing CPR:

Once CPR has been initiated, it can be discontinued when any one of the following occurs:

- 1) A "no code" or DNR order is confirmed
- 2) The patient has been transported to a health care facility and a physician at the facility determines that CPR is futile or should be stopped
- 3) A physician on scene or medical control for the service, based on information from members of the service on scene, determines that CPR is futile or should be stopped,
- 4) An out-of-hospital emergency care provider is following termination of CPR physician medical director approved protocols.

VIII. Refusal of Care

A. Adults

An adult is an individual 19 years old or older or who is or has been married (NEB REV STAT §43-2101). A competent adult can refuse medical services and/or transportation to a health care facility.

1. A legal guardian can consent to or refuse medical services and/or transportation to a health care facility for an incompetent adult.
2. A person appointed as a health Care Power of Attorney can consent or refuse consent for medical services and/or transportation to a health care facility for the incompetent adult named in the power of attorney.

B. Minors

A minor is an individual under 19 years of age that has never been married. A parent or legal guardian can consent or refuse consent on behalf of a minor, for medical services and/or transportation to a health care facility.

C. Documentation

All consents or refusals of consents for medical treatment and/or transportation must be documented in the patient care report. When possible these should be signed and dated by the patient or other individual authorized to give or refuse consent. All refusals to sign a consent or refusal of consent must be documented in the patient care report or other appropriate record(s).

IX. Transport/Intercepts:

- A. Patients should be transported as soon as possible to an appropriate medical facility. Generally, on-scene times for trauma patients should not exceed ten (10) minutes. Immediate transport with treatment en route is required for patients with significant trauma, unstable airways, or a patient with signs and symptoms of a Cerebrovascular accident (CVA)
- B. First Responder use, while encouraged, should not be used to replace EMT level and above providers whose training level is required for long term treatment and transport of patients.
- C. Tiered response, with an appropriate service is encouraged if assistance or level of care needs exist and can be met in a timely manner.

X. Communications:

To allow for regional or local variations, and needs, the provider may follow locally established and physician medical director approved communications policies and procedures.

Contact medical control as soon as feasible in accordance with local guidelines for on-scene or en route orders. For seriously injured or critically ill patients, give a brief initial report from the scene when possible with more detailed information given to medical control while en route.

1. When communicating with medical control or the receiving facility, a verbal report may include these essential elements:
 - a. Identify unit (If ALS staffed identify by “[town] medic ____”. If BLS staffed, identify by “[town] ambulance ____”.)
 - b. Patient’s age, gender
 - c. Patient’s chief complaint
 - d. Brief pertinent history of present illness or mechanism of injury (MOI)
 - e. Major past illnesses
 - f. Mental status
 - g. Baseline vital signs
 - h. Pertinent findings of the physical exam
 - i. Emergency medical care given
 - j. Patient response to treatment
 - k. Estimated time of arrival (ETA)
2. Advise the receiving facility of changes occurring in patient’s status en route.
3. Complete patient care report and provide a copy before leaving the receiving facility to assure continuity of patient care.

XI. After the Call:

- A. Notify dispatch when back in service. Clean, restock, and check over vehicle and equipment for next assignment.
- B. Consider having a Critical Incident Stress Debriefing (CISD) anytime rescuers and health care providers have been involved in a major incident, or one that produces adverse reaction.
- C. Remember the importance of patient confidentiality.

GENERAL PRINCIPLES

Airway and Oxygen

- A. An intact airway and adequate oxygenation is essential for all patients with medical or traumatic illnesses. Throughout this treatment protocol it is assumed that the Out-of-Hospital emergency care provider will maintain a patent airway and provide appropriate supplemental oxygenation.
 - 1. Adequate ventilations are defined as:
 - a. rate of 10-30
 - b. absence of shallow or labored effort
 - c. clear lung sounds
 - d. no or very little signs of distress
 - 2. Inadequate ventilations are defined as:
 - a. rate <10 or >30
 - b. a rate between 10-30 in the presence of:
 - shallow/labored respirations
 - OR
 - wheezes, wet sounds (crackles [rales] or bubbles)
 - OR
 - blue, gray or mottled skin
- B. Establish and maintain a secure airway/ventilation
 - 1. If ventilating adequately: nasal cannula O₂, 2-6 L/min. or Non-Rebreather Mask (NRB mask) at 10-15 L/min
 - 2. Maintain patent upper airway with jaw thrust, nasopharyngeal pharyngeal and/or oropharyngeal airway
 - 3. If not ventilating adequately: Assist with BVM and 100% O₂.
 - 4. If vital signs have not improved after initial oxygen, re-evaluate oxygen delivery and adjust accordingly.
 - 5. If pulse oximetry is used, adjust oxygen delivery devices to an oxygen saturation of 90% or above (goal is 100%) if possible.
 - 6. In case of cervical compromise, consider alternative techniques including use of lighted stylet, multi-lumen airway, or trauma ET intubation.
 - 7. Rapid Sequence Intubation (RSI), Needle cricothyrotomy, and surgical cricothyrotomy are advanced alternative techniques for airway management that require specialized training and authorization by the service program medical director. (See Appendix 1)

- C. Criteria for the use of age appropriate ventilation and advanced airway adjuncts:
1. Criteria for use of bag- valve-mask (BVM):
 - a. inadequate ventilation
 - b. rate <10 or >30; <20 in a pediatric patient with altered mental status
 - c. able to say only short phases/words before running out of breath
 - d. unconscious person with pale, cyanotic or gray color
 - e. irregular respirations
 - f. grunting in the pediatric patient
 2. Criteria for use of advanced airway management skills (multilumen airway, lighted stylet, LMA, oro or nasotracheal tube intubation)
 - a. unconscious patient who is apneic
 - b. patient who is unresponsive to painful stimuli
 - c. patient with no gag reflex or does not cough
 - d. inability of the patient to protect his/her own airway, i.e. cardiac arrest, unresponsive, etc.

Patient Assessment

- A. First Responders conduct a SIMPLE assessment as appropriate for the patient's condition. A Simple assessment includes assessing the presence or absence and quality of the ABC's, a determination of the patient's mental status, and a SAMPLE history. (As detailed in the DOT First Responder Curriculum)
- B. EMT-Basics EMT Options and EMT Intermediate 85s conduct a BASIC assessment as appropriate for the patient's condition. A Basic assessment includes performing an initial assessment, focused assessment, detailed assessment, and on going assessment. (As detailed in the DOT EMT-Basic Curriculum)
- C. EMT Intermediate 99s conduct an Advanced Physical Assessment as appropriate for the patient's condition. This assessment includes the Basic assessment as well a detailed assessment of lung sounds, abdomen, and the extremities. (As detailed in the 1999 DOT EMT- Intermediate Curriculum)
- D. Paramedics conduct an Comprehensive assessment as appropriate for the patient's condition. This assessment includes the Basic, Advanced Physical Assessments as well as a comprehensive assessment of body systems. (As detailed in the DOT EMT-Paramedic Curriculum)

IV Therapy

- A. If an advanced level intervention for an unstable patient requires IV access, the IV should be started as soon as feasible. For trauma patients, IV's should be started en route to the hospital, except when there is an unavoidable delay (long extrication, etc.).

- B. All IV insertions refer to peripheral IV's, including saline locks and intraosseous (IO) lines.
- C. Venous access can be achieved using either:
 - 1. Saline lock – used on patients who have stable vital signs and do not require volume replacement
 - 2. IV of Normal Saline (0.9% Sodium Chloride) or Lactated Ringers for IV fluid administration
- D. IV fluid administration is at the following rates:
 - 1. TKO – slow drip for patients that may need IV medication or fluid bolus
 - 2. Fluid Challenge – rapid 250-500 ml fluid bolus (Pediatric: 20 ml/kg)
 - 3. Maintain IV flow rate as ordered by physician/standing order
- E. Pre-existing Venous Access Devices (VAD) may be used in emergency situations
 - 1. Minidrip means IV administration set that delivers 60 gtts/ml
 - 2. Maxidrip means IV administration set that delivers 10, 15, or 20 gtts/ml

Patient Condition

- A. Unstable Patient – The patient as a whole must be assessed and no single parameter defines if the patient is unstable. Listed are guidelines for determining an unstable patient.
 - 1 Adult
 - a. Level of Consciousness – decreased or decreasing
 - b. Blood Pressure <90 in the adult patient
 - c. Chest pain
 - d. Dyspnea
 - e. Signs and symptoms of uncompensated shock
 - f. Patient does not respond to treatment(s) and continues to worsen
 - 2 Pediatric
 - a. Level of Consciousness – Decreased for the age of the patient, a lethargic, limp young child or infant should be considered unstable
 - b. Cool, clammy, pale skin
 - c. Signs of dehydration
 - 1. decreased urine output
 - 2. increased pulse rate for age group
 - 3. dry mucus membranes
 - 4. sunken eyes/fontanelle
 - d. Signs and symptoms shock progressing rapidly

- e. Signs and symptoms of uncompensated shock
 - f. Patient does not respond to treatment(s) and continues to worsen
- B. Stable Patient – As with the unstable patient the patient as a whole must be assessed. Listed are guidelines for determining a stable patient
 - 1 Adult and Pediatric
 - a. Awake alert
 - b. Blood pressure – normal range
 - c. Skin – normal
 - d. Patient responds to treatments and improves

Transport

- A. Non Emergent – patient transport without the use of emergency lights and siren. Most patients can be transport non-emergent to the hospital
 - a. Patient is stable
 - b. Weather/road/traffic conditions prevent safe emergent driving
 - c. Use of lights/siren would agitate a patient or exacerbate the patients condition. Examples:
 - i. Chest Pain may be made worse
 - ii. Seizures may be induced
 - iii. Airway occlusion may be cause in the epiglottitis patient
- B. Emergent Transport – patient transport with the use of lights and siren as needed
 - a. The Patient is unstable

ADULT CARDIOVASCULAR EMERGENCIES**Cardiopulmonary Arrest****First Responder**

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Confirm Unresponsiveness
AED	Attach AED Do Not Delay AED for CPR ** SEE AED Protocol***
Airway	Establish an Airway
Breathing	Ventilate with Bag Valve Mask Attached to O2 OR Pocket Mask
Circulation	Confirm Patient Pulseless and Begin Chest Compressions
Prepare	Package for Transport

EMT

Airway	Insert an Oral Airway
Transport	Emergent
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Airway	Insert Medical Director Approved Advanced Airway Device (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access

EMT Intermediate 99

Cardiac Monitor/ Defib	Attach Monitor/Defib Unit Interpret Rhythm
ACLS	Follow Appropriate <u>Adult Cardiac Arrest Algorithm</u>

Paramedic

ACLS	Follow Appropriate <u>Adult Cardiac Arrest Algorithm</u>
------	--

Note: If the patient regains a pulse see the Adult Post Cardiac Arrest – Return of Pulse protocol

Adult Cardiac Arrest Algorithm V-Fib

EMT-Intermediate 99

Shock	3 Shocks Monophasic or Biphasic
Airway	Establish an Airway with an Advanced Airway Device Endotracheal Intubation Preferred
Breathing	Ventilation with Bag Valve Mask Attached to O ²
Circulation	Chest Compression and Establish Peripheral IV Access
Medication	Epinephrine 1mg Every 3 – 5 Minutes
Shock	Shock
Intervention Sequence	Drug Shock Repeat
Medication Considerations	Lidocaine 1-1.5 mg/kg

If IV access is delayed or can not be obtained Epi may be given via the ET tube

If the patient's cardiac rhythm changes see appropriate protocol for that rhythm.
If patient develops a pulse see the Post Cardiac Arrest protocol.

Paramedic

Medication Considerations	Vasopressin in lieu of First Dose of Epinephrine 40U Single Dose if No Response in 10 to 20 Minutes May Return to Epinephrine
	Amiodarone 300mg 1 st dose 150mg 2 nd dose
	Magnesium Sulfate 1-2g For Torsades de Pointes
	Procainamide Infuse 50mg/min to max of 17mg/kg Do Not Use Both Procainamide and Amiodarone on the Same Patient
	Sodium Bicarbonate 1mEq/kg

Adult Cardiac Arrest Algorithm PEA

EMT-Intermediate 99

Confirm	Cardiac Rhythm
Airway	Establish an Airway with an Advanced Airway Device Endotracheal Intubation Preferred
Breathing	Ventilation with Bag Valve Mask Attached to O ²
Circulation	Chest Compression and Establish Peripheral IV Access
Medication	Epinephrine 1mg Every 3 – 5 Minutes
Intervention Sequence	Drug Evaluate for Change Repeat
Medication Considerations	Atropine 1mg (If Rate Slow) Repeat Atropine Every 3-5 Minutes to Max Dose .04mg/kg
Consider	Consider Causes of PEA Hypovolemia Consider Fluid Boluses Tension Pneumothorax Consider Needle Decompression Hypothermia Consider Warming Patient

If IV access is delayed or can not be obtained Epi and Atropine may be given via the ET tube

If the patient's cardiac rhythm changes see appropriate protocol for that rhythm. If patient develops a pulse see the Post Cardiac Arrest protocol.

Paramedic

Consider	Consider Causes of PEA Acidosis Consider Sodium Bicarbonate Tricyclic Overdose Consider Sodium Bicarbonate Calcium Channel Blocker Overdose Consider Calcium Chloride
Medication Doses	Sodium Bicarbonate 1mEq/Kg IVP Calcium Chloride 500 to 1000mg Slow IVP

Adult Cardiac Arrest Algorithm Asystole

EMT-Intermediate 99

Confirm	Cardiac Rhythm
Airway	Establish an Airway with an Advanced Airway Device Endotracheal Intubation Preferred
Breathing	Ventilation with Bag Valve Mask attached to O2
Circulation	Chest Compression and Establish Peripheral IV Access
Medication	Epinephrine 1mg Every 3 – 5 Minutes
Intervention Sequence	Drug Evaluate for change Repeat
Medication Considerations	Atropine 1mg Repeat Atropine Every 3-5 Minutes to Max Dose .04mg/kg
Consider	Consider Causes Asystole Hypothermia Consider Warming Patient

If IV access is delayed
or can not be obtained
Epi and Atropine may
given via the ET tube

If the patient's cardiac
rhythm changes see
appropriate protocol
for that rhythm.
If patient develops a
pulse see the Post
Cardiac Arrest

Paramedic

Consider	Transcutaneous Pacing
Consider	Consider causes Acidosis Consider Sodium Bicarbonate Tricyclic Overdose Consider Sodium Bicarbonate Hyperkalemia Consider Sodium Bicarbonate Calcium Channel Blocker Overdose Consider Calcium Chloride
Medication Doses	Sodium Bicarbonate 1mEq/Kg IVP Calcium Chloride 500 to 1000mg Slow IVP

AED Protocol

ALL LEVELS

Confirm	Patient is Pulseless		
AED	Attach AED Pads and Turn ON Do Not Delay AED for CPR		
Analyze	Push Analyze Button		
Follow Voice Prompt	Push to Shock	OR	No Shock Advised Check Pulse If None Do 1 minute of CPR
Analyze	Push Analyze Button		Push Analyze Button
Follow Voice Prompt	Push to Shock	OR	No Shock Advised Check Pulse If None Do 1 Minute of CPR Package for Transport Transport if EMT or Higher Level
Analyze	Push Analyze Button		Push Analyze Button
Follow Voice Prompt	Push to Shock	OR	No Shock Advised
Pulse Check	Check Pulse If none do 1 minute of CPR		
Analyze	Push Analyze Button		Push Analyze Button
Follow Voice Prompt	Push to Shock	OR	No Shock Advised Check Pulse If None Do 1 Minute of CPR Package for Transport Transport if EMT or Higher Level
Repeat	Analyze and Shock Three More Times		
Pulse Check	Check Pulse if None Package for Transport Transport if EMT or Higher Level		
Analyze/Shock	Each 3-5 Minutes Push Analyze if Shock Advised Shock Up to Three Times and then Continue Transport		

Considerations:

If the “No Shock Advised” prompt is heard after three consecutive analyze steps continue CPR and package for transport and for EMT and higher level begin transport.

If No shock advised and patient has return of pulse see Post Cardiac Arrest – Return of Pulse protocol.

Post Cardiac Arrest – Return of Pulse

With public access to AEDs the OOH care provider may respond to a person who has been successfully defibrillated
OR the OOH provider may successfully defibrillate a cardiac arrest victim.

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Confirm Unresponsiveness
Airway	Establish an Airway
Breathing	Assist Ventilations with Bag Valve Mask
Circulation	Confirm Pulse Present, Recheck Often
Assess	Conduct Simple Patient Assessment

EMT

Airway	Insert Oral Airway
Assess	Perform Basic Patient Assessment
Consider	ALS Intercept

EMT Options / EMT Intermediate 85

Airway	Insert Medical Director Approved Advanced Airway Device if Indicated (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access
Consider	250cc Fluid Bolus

EMT Intermediate 99

Assess	Perform Advanced Assessment
Cardiac Monitor	Determine Cardiac Rhythm
Consider	Lidocaine 1mg/kg bolus if rate >60 and presence of ventricular ectopy Lidocaine infusion 1 – 4 mg/min
Dysrhythmias	Treat with Appropriate <u>Advanced Cardiac Dysrhythmia</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Consider	12 Lead EKG
Dysrhythmias	Treat with Appropriate <u>Advanced Cardiac Dysrhythmia</u> Protocol
Consider	Dopamine Infusion 5 to 20 mcg/kg/min for Hypotension

Discontinue CPR

Situations may occur where CPR has been initiated on an obviously deceased patient prior to the arrival of out-of-hospital emergency care providers.

All certification levels

If the following criteria have been met, the out-of hospital emergency care providers may discontinue CPR or may choose not to initiate CPR:

1. No pulse; AND
2. No spontaneous respirations; AND
3. Pupils fixed and dilated; AND
4. One or more of the following:
 - A. Rigor mortis;
 - B. Decapitation;
 - C. Decomposition;
 - D. Dependent lividity;
 - E. Traumatic cardiopulmonary arrest with injuries incompatible with life (i.e. massive blood loss, displacement of brain tissue);
 - F. Valid DNR form; or
 - G. Physician authorization;
5. Determination of the patient's cardiac rhythm is not required

NOTE:

Patients in whom hypothermia may be a significant component of their arrested state should receive resuscitative efforts until body core temperature is >35 degrees centigrade.

General Cardiac Dysrhythmia

Dysrhythmia-An abnormal heart rate and/or rhythm

The First Responder, EMT, EMT with Option, and EMT-I 85 will only detect by evaluation of the pulse whether the pulse is fast, slow, or normal rate and if it is regular or irregular. EMT-I 99 and Paramedic will interpret the Cardiac Monitor strip to determine the type of cardiac rhythm.

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Painful, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Assess Pulse Rate, Rhythm, and Quality Vital Signs
Assess	Conduct Simple Patient Assessment
Prepare	Prepare Patient for Transport

EMT

Assess	Perform a Basic Assessment
Determine	Patient Stable or Unstable
Transport	Non-emergent Transport Unless Patient Unstable
Consider	ALS Intercept

EMT Options / EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT Intermediate 99

Assess	Perform Advanced Assessment
Cardiac Monitor	Determine Cardiac Rhythm
ACLS	Follow Appropriate <u>Advanced Cardiac Dysrhythmia</u> Protocol

Paramedic

Consider	12 Lead ECG
Assess	Perform Comprehensive Assessment
ACLS	Follow Appropriate <u>Advanced Cardiac Dysrhythmia</u> Protocol

Advanced Cardiac Dysrhythmia

This Protocol applies to EMT-I 99 and Paramedics and assumes the General Cardiac Dysrhythmia protocol has been followed

The EMT –I99 and Paramedic must determine if the patient is stable or unstable

For the stable patient tolerating the cardiac rhythm; May require monitoring and supportive care without a medication or electrical intervention.

For the unstable patient the EMT-I 99 or Paramedic; May have to consider a more aggressive treatment plan based on the patient condition and how rapidly a medication may be delivered verses an electric therapy can be performed.

The Paramedic may choose a medication intervention from either the EMT-I 99 or the Paramedic sections.

Ventricular Ectopy PVC's , Couplets, Bi and Trigeminy Heart Rate Above 60

EMT Intermediate 99

Consider	Lidocaine 1mg/kg Bolus
If Dysrhythmia resolves	Lidocaine Infusion 1-4mg/min

Paramedic

Consider	Amiodarone 150mg Over 10 Minutes
----------	----------------------------------

Ventricular Tachycardia

	Stable	Unstable
EMT Intermediate 99		
Consider	12lead ECG	
Consider	Lidocaine 1mg/kg	Synchronized Cardioversion Premedicate if Possible Diazepam 2-5mg
If Dysrhythmia Resolves	Lidocaine Infusion 1-4mg/min	Lidocaine 1mg/kg Bolus Followed by Lidocaine Infusion 1-4mg/min

Paramedic

Consider	Amiodarone 150mg Over 10 Minutes or Procainamine 20mg/min until resolved Or Magnesium Sulfate 1-2g (if Polymorphic)	Alternate Pre-medications Midazolam 2-4mg or Lorazepam 2 -4 mg
Consider	Infusion of Anti-Arhythmic Agent that Controlled the Dysrhythmia	

Advanced Cardiac Dysrhythmia Continued**Atrial Tachycardias**

PSVT, Atrial Fib, Atrial Flutter

	Stable	Unstable
EMT Intermediate 99		
A-Fib A Flutter		
Consider	Confirm Rhythm	Synchronized Cardioversion Premedicate if Possible Diazepam 2-5mg
PSVT		
Consider	Adenosine Rapid IVP 6mg then 12mg	Synchronized Cardioversion Premedicate if Possible Diazepam 2 to 4 mg
Paramedic		
A-Fib A Flutter		
Consider	12 Lead	Alternate Pre-medications Midazolam 2-4mg or Lorazepam 2-4mg
PSVT		
Consider	12 Lead	
Consider	Verapamil 2.5-5mg or Diltiazem .25mg/kg	Alternate Pre-medications Midazolam 2 4mg or Lorazepam 2-4mg

Must confirm that A-Fib A-Flutter is new onset before cardioversion

Bradycardia

	Stable	Unstable
EMT-Intermediate 99		
Consider	12 Lead	Atropine .5-1mg ¹
Consider	Have Pacer Standing By	Transcutaneous Pacing ² Premedicate if Possible Diazepam 2 to 4 mg
Paramedic		
Consider	If Second or Third Degree Block Attach Pacer Pads	Alternate Pre-medications Midazolam 2 4mg or or Lorazepam 2-4mg
Consider		Dopamine 5 -20mcg/kg/min or Epinephrine 2-10mcg/min or Isoproterenol 2-10mcg/min

1. Atropine is not effective in Second Degree Type II and new Third Degree Heart Blocks.
Atropine is not effective for denervated transplanted hearts
2. Do not delay pacing for IV or pre-medication if patient is deteriorating.

Chest Pain**First Responder**Scene Safety –
BSI

SAFETY FIRST

Level of
Conscious

Alert, Verbal, Pain, or Unresponsive

Airway

Monitor Airway

Breathing

Administer Oxygen

Circulation

Vital Signs, Skin Color/Temp

Assess

Conduct Simple Patient Assessment

Consider

Aspirin – Two to Four – 81mg (Baby Aspirin) Chewed and Swallowed
Only if the OOH Provider has been Trained and Approved by the Service's Medical Director**EMT**

Assess

Conduct Basic Patient Assessment

Consider

May Assist Patient with Taking his/her Own Nitroglycerin .4mg Tablet or Spray
Sublingually
May Repeat up to Three Times if BP Remains >100 Systolic

Transport

Non-emergent Transport unless patient becomes unstable

Consider

ALS Intercept

EMT Options/ EMT Intermediate 85

IV

Establish Peripheral IV Access

EMT-Intermediate 99

Assess

Perform Advanced Physical Assessment

Cardiac
Monitor

Attach Cardiac Monitor, Interpret ECG

Consider

Nitroglycerin – One 0.4mg Tablet or Spray Sublingually
Repeat Every 5 Minutes if Chest Pain Continues AND BP Remains >100 Systolic

Morphine

Administer 2 – 5 mg IV
May Repeat PRN Until Pain Relieved AND Blood Pressure Remains >100 systolic

Dysrhythmia

*SEE Cardiac Dysrhythmia Protocol**Paramedic**

Assess

Perform Comprehensive Assessment

Consider

12 Lead

Dysrhythmia

*See Cardiac Dysrhythmia Protocol

Alternate

If allergic to Morphine may use Fentanyl 25mcg to 100mcg IV

Pulmonary Edema

Dyspnea in the presence of diminished lung sounds, wheezes, rales, or frothy sputum with a BP that is hypertensive or within normal limits.

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Consider	Assisting Ventilations
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non-emergent Transport Unless Patient Becomes Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Consider	Nitroglycerin – One 0.4mg Tablet or Spray Sublingually if BP >100 Systolic
Consider	Furosemide 40-80mg IV
Consider	Administer Morphine 2 – 4 mg IV
Consider	Bronchodilator Medication by Nebulizer *See <u>Bronchodilator</u> Protocol
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Consider	12 Lead
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Cardiogenic Shock

**Dyspnea in the presence of diminished lung sounds, wheezes, rales, or frothy sputum
with a BP that is hypotensive**

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Consider	Assisting Ventilations
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Emergent Transport
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Consider	Bronchodilator Medication by Nebulizer *See <u>Bronchodilator</u> Protocol
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol
Consider	Dopamine 5 to 20mcg/kg/min infusion
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Difficulty Breathing
Acute Allergic Reaction / Anaphylaxis

Difficulty Breathing in the presence of urticaria, wheezing and /or contact with a known allergen

First Responder

Scene Safety –

BSI SAFETY FIRST

Level of

Conscious Alert, Verbal, Pain, or Unresponsive

Airway

Monitor Airway

Breathing

Administer Oxygen, Consider Assisting Ventilations With BVM

Circulation

Vital Signs, Skin Color/Temp

Assess

Conduct a Simple Patient Assessment

First Responder AND EMT

Consider

Epinephrine Auto Injector (EPI PEN)

Only if the OOH Provider has been Trained and Approved by the Service's Medical Director Guidelines; Patient able to speak one-two word phrases

Low/falling oxygen saturations even with O2 administration

Diminished to absent lung sounds

Decreasing LOC

Retractions

Pale or cyanotic skin

EMT

Assess

Conduct Basic Patient Assessment

Consider

Assist Patient with His/Her Metered Dose Inhaler (MDI)

Transport

Emergent Transport Unless Patient is Stable

Consider

ALS Intercept

EMT Options/ EMT Intermediate 85

IV

Establish Peripheral IV Access titrate to blood pressure

Continued next page

Acute Allergic Reaction / Anaphylaxis

Continued

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Epinephrine 1:1000 .3-.5mg SubQ May Repeat Every 5 Minutes if Severe Respiratory Symptoms are Not Resolved or Epinephrine 1:10,000 .1-.2mg IV For BP < 70 Systolic May Repeat Every 5 Minutes if Severe Respiratory Symptoms are Not Resolved
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Consider	Bronchodilator Medication by Nebulizer ***See <u>Bronchodilator</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Diphenhydramine (Benadryl), 25 – 50 mg Slow IVP over 1-3 Minutes
Consider	Solu-medrol 125mg IVP
Consider	Dopamine for BP < 70 Systolic 5 to 20mcg/kg/min Infusion
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Asthma**Difficulty Breathing in the presence of wheezing with a history of asthma or irritant exposure****First Responder**

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen - Consider Assisting Ventilations with BVM
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

First Responder AND EMT

Consider	Epinephrine Auto Injector (EPI PEN) if Impending Respiratory Collapse Only if the OOH Provider has been Trained and Approved by the Service's Medical Director Guidelines; Patient able to speak one-two word phrases Low/falling oxygen saturations even with O2 administration Diminished to absent lung sounds Decreasing LOC Retractions Pale or cyanotic skin
----------	---

EMT

Assess	Conduct Basic Patient Assessment
Consider	Assist Patient with His/Her Metered Dose Inhaler (MDI)
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Bronchodilator Medication by nebulizer *See <u>Bronchodilator</u> Protocol
Consider	Epinephrine 1:1,000 .3-.5mg SubQ
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

COPD
Emphysema or Chronic Bronchitis
Difficulty Breathing in the presence of wheezing and/or rhonchi and history of COPD

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen - Consider Assisting Ventilations with BVM
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Consider	Assist Patient with His/Her Metered Dose Inhaler (MDI)
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Bronchodilator Medication by nebulizer *See <u>Bronchodilator</u> Protocol
Consider	Epinephrine 1:1,000 .3-.5mg SubQ
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess	Perform Comprehensive Assessment
Consider	CPAP
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Respiratory Infection

Difficulty Breathing in the presence of or suspected presence of Respiratory Infection

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Consider	Assist patient with His/Her Metered Dose Inhaler (MDI)
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Bronchodilator Medication by nebulizer *See <u>Bronchodilator</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
--------	----------------------------------

***Note: Clean unit and equipment with an appropriate disinfectant after call

Adult Epiglottitis

Difficulty breathing in the presence of stridor.

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

Epiglottitis may cause the patient airway to become occluded completely if the patient is agitated.

EMT

Assess	Conduct Basic Patient Assessment
Airway	DO NOT INSERT ORAL OR NASAL AIRWAY
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access EXTREME CAUTION MUST BE EXERCISED
----	---

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG

Epiglottitis may cause the patient airway to become occluded completely if the patient is agitated.

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Cricothyrotomy Needle or Surgical For Occluded Airway

ACUTE MEDICAL EMERGENCIES

Upper Airway Obstruction

First Responder

Scene Safety –
BSI

SAFETY FIRST

Level of
Conscious

Alert, Verbal, Pain, or Unresponsive

Airway

Attempt to Relieve Obstruction Using AHA Guidelines For Obstructed Airway

Continue

Obstruction not Cleared
Attempts to relieve
Obstruction

Obstruction Cleared

Breathing

Administer Oxygen
Consider Assisting Ventilations
with BVM

Circulation

Vital Signs, Skin Color/Temp

Assess

Conduct a Simple Patient Assessment

EMT

Transport

Emergent
Continue Attempts to Relieve
Obstruction

Non-emergent Transport if Patient Stable

Assess

Conduct Basic Patient Assessment

Consider

ALS Intercept

ALS Intercept

EMT Options/ EMT Intermediate 85

Airway

Attempt to Visualize Obstruction with
Laryngoscope and Remove with McGill
Forceps. IF TRAINED

Consider Advance Airway if Patient
LOC Remains Decreased and No Gag Reflex

IV

Establish Peripheral IV Access
Do Not Delay Transport

Establish Peripheral IV Access

EMT-Intermediate 99

Assess

Perform Advanced Physical Assessment

Cardiac Monitor

Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess

Perform Comprehensive Assessment

Consider

Cricothyrotomy needle or surgical

Rapid Sequence Intubation (RSI)

*See RSI Protocol

Non-Traumatic Altered or Decreased Level of Consciousness
Diabetes Mellitus – Hypoglycemia
Altered Mental Status with History of Diabetes Mellitus (Hypoglycemia)

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Consider	Oral glucose ONLY if Patient's Airway can be Maintained
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Glucometer	Obtain Glucose Reading (Reading >200 *See <u>Hyperglycemia</u> Protocol)
IV	Establish Peripheral IV Access

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Dextrose 50% 25g IVP

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Thiamine 100mg IV Prior to Dextrose 50%
Consider	Glucagon, 0.5-1.0 mg IM or Subcutaneously If Unable to Obtain IV

Hyperglycemia

Hyperglycemia with or without known history of Diabetes Mellitus

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Glucometer	Obtain Glucose Reading
Consider	Medical Director Approved Advanced Airway Device if Indicated (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access Run Wide Open, Monitor for Fluid Overload Consider 2 nd IV in Presence of Profound Dehydration

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Consider	Narcan .4mg to 2mg Only if Suspected Drug Overdose

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Altered Mental Status**Altered Mental Status in the Absence of Exposure and No History of Diabetes Mellitus****First Responder**

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Emergent Unless Patient LOC Improves with Glucose
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Glucometer	Obtain Glucose Reading (Reading >200 *See <u>Hyperglycemia</u> Protocol)
Consider	Oral Glucose if Indicated by Glucometer reading of <80 <u>AND</u> if Patient's Airway can be Maintained
Consider	Inserting Medical Director Approved Advanced Airway Device (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Dextrose 50% 25g IVP if Indicated by Glucometer Reading <80
Consider	Narcan .4mg to 2mg IV if Drug Overdose is Suspected

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Thiamine 100mg IV Prior to Dextrose 50%
Consider	Glucagon, 0.5-1.0 mg IM or Subcutaneously If Unable to Obtain IV AND Indicated by Glucometer Reading
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Cerebrovascular Accident (CVA)

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non emergent Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Consider	Inserting Medical Director Approved Advanced Airway Device if Indicated (Multi-lumen Airway, LMA, ET)
Glucometer	IF LOC Decreased, Obtain Glucose Reading and Use other Appropriate Protocol if Abnormal
IV	Establish Peripheral IV Access

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Seizure Disorder

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment
Prepare	Package Patient for Transport

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non –Emergent Transport Unless Seizures Continue
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Glucometer	Obtain Glucose Reading and Use Other Appropriate Protocol if Abnormal
IV	Establish Peripheral IV Access

EMT-Intermediate99

Assess	Perform Advanced Physical Assessment
Consider	Possible Causes of Seizure
Consider	Diazepam 5-10mg IV for Recurrent or Prolonged Seizures
Consider	Cardiac Monitoring

Paramedic

Assess	Perform Comprehensive Assessment
Alternate Medication	May Consider Lorazepam 2-4mg alternate to Diazepam

Hypothermia

Lowered skin Temperature with Altered Mental Status

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Confirm Unresponsiveness
Airway	Establish an Airway
Breathing	Administer Warmed Oxygen if Possible If Not Breathing or Respiratory Compromised Ventilate with Bag Valve Mask Attached to O2 or Pocket Mask
Circulation	Confirm Pulse If Pulseless Attach AED Do Not Delay AED for CPR * See AED Protocol Begin Chest Compressions
Avoid	Rough Handling/Movement
Warm	Remove Wet Clothing Remove Patient From Cold Warm Body Core – Heat packs to Groin and Axillary Areas
Prepare	Package for Transport

EMT

Airway	Insert an Oral Airway if Indicated
Temp	Obtain Body Temperature

Consider ALS Intercept

EMT Options / EMT Intermediate 85

Glucometer	Obtain Glucose Reading use Other Appropriate Protocol if Abnormal
Airway	Insert Medical Director Approved Advanced Airway Device If Indicated (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access **WARM IV FLUID**

EMT Intermediate 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
ACLS	Follow Appropriate Cardiac Arrest Algorithm *NOTE ACLS Medications Should be Avoided Until Patient Warmed

Paramedic

Assess	Perform Comprehensive Assessment
ACLS	Follow Appropriate Cardiac Arrest Algorithm *NOTE ACLS Medications Should be Avoided Until Patient Warmed

Hyperthermia

Elevated skin Temperature with altered Mental Status

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment
Cool	Remove Layers of Clothes, Place Patient in Cool Environment, Wrap Patient in Moist Sheets
Prepare	Package Patient for Transport

EMT

Assess	Conduct Basic Patient Assessment
Airway	Insert a Oral Airway if Indicated
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Glucometer	Obtain glucose reading use other appropriate protocol if abnormal
Airway	Insert Medical Director Approved Advanced Airway Device if Indicated (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Consider	Valium 5-10mg IV for Seizures
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol

Non-Traumatic Abdominal Pain

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Consider	Cause of abdominal pain
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Pain Management *See <u>Pain Management</u> Protocol
Consider	Cardiac Monitoring

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Pain Management **See <u>Pain Management</u> Protocol

Upper and Lower Gastrointestinal Bleeding

Coffee ground emesis, Tarry black stools, with or without abdominal pain

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IVs Titrate to BP Consider 2 nd IV for Additional Fluid Challenge
----	--

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Pain Management **See <u>Pain Management</u> Protocol
Consider	Cardiac monitoring

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Pain Management **See <u>Pain Management</u> Protocol

Hypotension in the Absence of Trauma

Hypovolemic Shock

Shock present when pulse greater than 120 and systolic BP less than 100 mmHg in a previously normotensive patient; OR systolic drops 40-50 mmHg in a previously hypertensive patient, especially if accompanied by pale, clammy skin, decreased level of consciousness, and poor capillary refill.

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access, Titrate to BP Consider 2 nd IV for Additional Fluid Challenge
----	---

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Cardiac Monitoring

Paramedic

Assess	Perform Comprehensive Assessment
--------	----------------------------------

OBSTETRICS and GYNECOLOGIC EMERGENCIES**Imminent Delivery with History of Pregnancy, a Palpable Uterus and Contractions****First Responder**

Scene Safety –

BSI SAFETY FIRST

Level of

Conscious Alert, Verbal, Pain, or Unresponsive

Airway

Monitor Airway

Breathing

Administer Oxygen

Circulation

Vital Signs, Skin Color/Temp

Assess

Conduct a Simple Patient Assessment

Prepare

Mother for Delivery if Crowning

Delivery

Use OB Kit and Deliver Infant

*See Newborn Care Protocol

Post

Partum Care

Allow placenta to deliver naturally. Massage top of uterus, put baby to breast.

Bring all tissue passed to the hospital. **DO NOT** forcibly extract any tissue.

Place OB Pad

*See Newborn Care Protocol

Monitor

Mother for severe postpartum bleeding

Control Post Partum Bleeding by Massaging the Top of Uterus,
and Put Baby to Breast

Do Not Pack Anything in the Vagina

EMT

Assess

Conduct Basic Patient Assessment

Transport

Non –Emergent Transport

Consider

ALS Intercept

EMT Options/ EMT Intermediate 85

IV

Establish Peripheral IV Access

EMT-Intermediate 99

Assess

Perform Advanced Physical Assessment

Paramedic

Assess

Perform Comprehensive Assessment

Newborn Care

First Responder

Scene Safety – BSI	SAFETY FIRST
Deliver	Support Head as it Passes From Birth Canal
Airway	Once the Head is Delivered, Suction Mouth Then Nose with Bulb Syringe EXAM Face/Head for Meconium Stained Fluid Suction Mouth, Pharynx, Nose until Clear
Cord	After the Delivery Keeping Baby at Level of Perineum, and Once Cord Stops Pulsating Double Clamp Cord, 6-12 inches from Baby and Cut Between Clamps
Dry/Warm	Once Fully Delivered, Dry and Wrap the Newborn
Breathing	Administer Blow by Oxygen Spontaneous Respirations Absent or <30, Slap or Flick the Soles of the Infant's Feet or Rub the Newborn's Back No Change in 5 Seconds Begin BVM Respirations Continue to Warm and Stimulate
Circulation	Pulse Rate < 100 BVM Respirations Pulse 60 to 80 and Not Rapidly Increasing Begin Chest Compressions Pulse < 60 Begin Chest Compressions
Assess	Assess Patient Each 5 Seconds for Changes Discontinue Chest Compression if Pulse Increases to 100 or Greater AND Maintains Discontinue BVM Resps. Once Spontaneous Breathing is >30 and Maintains

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non –Emergent Transport if Stable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Consider	Endotracheal Intubation <u>Only</u> Meconium – Intubate Suction Repeat Until Clear, Intubate With Clean Tube
----------	---

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	IV Access

Paramedic

Assess	Perform Comprehensive Assessment
--------	----------------------------------

Newborns benefit from rapid assessment and treatment of the ABC's and warming/stimulation. Rarely, advanced providers will need medication therapy. **EMT I99s and Paramedic see the Pediatric Cardiac Arrest and Dysrhythmia protocols for further guidance.**

Birth Complications**Arm or Leg Presentation, Prolapsed Cord, Significant Hemorrhage****First Responder**

Scene Safety –
BSI

SAFETY FIRST

Level of
Conscious

Alert, Verbal, Pain, or Unresponsive

Airway

Monitor Airway

Breathing

Administer Oxygen

Circulation

Vital Signs, Skin Color/Temp

Assess

Conduct a Simple Patient Assessment

Complications

Prolapsed Cord:

Place Patient on Back and Elevate the Hips – Consider Knees to Chest Position

Place Sterile-Gloved Index and Middle Fingers into the Vagina,

Push Infant Up to Relieve Pressure on Cord

Check Cord for Pulse.

Coach Mother to Breath Through Contractions and NOT to Push/Bear Down

Breech Delivery and Unable to Deliver Head:

Place Gloved Hand in the Vagina with Palm Towards Baby's Face

Form a V on Either Side of the Baby's Nose/ Mouth to Form Air Passage to Nose/ Mouth

Coach Mother to Breath Through Contractions and NOT to Push/Bear Down

Arm or Leg Presentation:

Place Patient on Back and Elevate the Hips – Consider Knees to Chest Position

Coach Mother to Breath Through Contractions and NOT to Push/Bear Down

If Significant Hemorrhage:

Place External Dressings, Monitor Bleeding and Elevate Hips

Coach Mother to Breath Through Contraction and NOT to Push/Bear Down

EMT

Assess

Conduct Basic Patient Assessment

Transport

Consider Emergent Transport

Consider

ALS Intercept

EMT Options/ EMT Intermediate 85

IV

Establish peripheral IV access

EMT-Intermediate 99

Assess

Perform Advanced Physical Assessment

Paramedic

Assess

Perform Comprehensive Assessment

Hypertensive Disorders of Pregnancy Toxemia of Pregnancy/Eclampsia

First Responder

Scene Safety- BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs
Assess	Conduct a Simple Patient Assessment
Position	Move Patient onto LEFT Side

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non-Emergent Unless Patient Becomes Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
--------	--------------------------------------

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Administer Magnesium Sulfate 2 - 6gm, Diluted to 25%, Slow IVP Over 3-5 Minutes, May Repeat Once After 5 Minutes.

Vaginal Bleeding with or without Gynecological Pain**First Responder**

Scene Safety- BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non-emergent Unless Patient Becomes Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
Consider	Pain Management **See Pain Management Protocol**

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Pain Management **See pain Management Protocol**

ACUTE TRAUMATIC EMERGENCIES
General Trauma Management– Priorities for Treatment

1. Body Substance Isolation and Scene Safety
2. Airway Management, Oxygen Administration, Vital Signs
3. Control the cervical spine. Assume cervical spine injury is present in any patient who has sustained trauma with:
 - a. Neurological deficit
 - b. Neck pain or tenderness with palpation
 - c. Altered mental status
 - d. Presence of a distracting injury
 - e. Any condition which may mask c-spine discomfort (i.e. recreational drug use, ETOH)
 - f. Any significant mechanism of injury
4. Remove all motor vehicle helmets to avoid airway management problems according to American College of Surgeons guidelines.
5. Football helmets should not be removed when shoulder pads are in place.
6. Control hemorrhage through
 - a. Direct pressure
 - b. Elevate effected extremity
 - c. Pressure dressing
 - d. Pressure points
 - e. Tourniquet (as a last resort), record time placed
7. Treat hypovolemic shock: Assume shock present when pulse greater than 120 and/or systolic BP less than 100 in a previously normotensive patient, especially if accompanied by pale clammy skin, decreased level of consciousness, and capillary refill > 2 sec.
8. Fractures/Dislocations – General Principles
 - a. Check and record peripheral pulses and neurological status before and after manipulating or splinting fractures.
 - b. Apply gentle in-line traction to fractures with the exception of dislocations or fractures involving joints (especially the elbow).
 - c. May straighten severely injured angulated fractures of extremities with exception of those involving knee or elbow (except if neurovascular bundle already compromised).
 - d. May use traction splint with open or closed femur fractures.
 - e. Immobilize fractures, including joint above and below site of fracture. DO NOT use inflatable splints for fractures of the humerus or the femur.
The PASG may be used to stabilize any fractures of the pelvis or lower extremities
9. DO NOT remove any impaled object unless obstructing airway
10. Pain Management should be consider once any life threatening injuries are treated AND vital sign indicate the patient is stable enough to tolerate the medication.
11. Consider 2 large bore IVs for all significant trauma. All IVs are to be titrated to vital signs

Trauma System

1. Evaluate dispatch information, the scene, mechanism of injury, patient presentation, distance to trauma center
 - a. Consider Helicopter Air Ambulance Standby (helicopter stays at it base on alert)
 - i. Reports of Penetrating and /or Significant Blunt Trauma to head, neck or torso
 - ii. Reports of entrapment
 - iii. Reports of burns and/or toxic inhalation injury
 - iv. Reports of decreased or loss of consciousness
 - v. Distance/ time from the scene to a designated trauma center exceeds 30 minutes
 - vi. Reports of a multiple patient incident
 - b. Consider Helicopter Air Ambulance response if
 - i. Patient condition indicates shock
 - ii. Confirmation of Penetrating and/or Significant Blunt Trauma to head, neck or torso
 - iii. Confirmation of amputated extremities above fingers/toes, and/or multiple long bone fractures
 - iv. Confirmation of burns >20% BSA or face /airway burns
 - v. Confirmation of a toxic inhalation injury with dyspnea
 - vi. Confirmation of entrapment with extrication/rescue needed
 - vii. Confirmation of decreased or loss of consciousness
 - viii. Confirmation of a multiple patient incident
 - ix. Location of the incident may allow the helicopter to make it first on a scene or at the same time as ground ambulance.
 - c. Additional Considerations in the request for Helicopter Air Ambulance
 - i. Your geographical distance from the Helicopter Air Ambulance, the local hospital (local trauma center), and regional trauma center.
 - ii. Time will be saved in delivering patient to a Trauma Center
 - iii. ALS level of care being delivered to the patient more timely
 - iv. Do not delay transport, consider an intercept with the helicopter if that can be done safely
 - v. Helicopter Air Ambulance may divert to prearranged landing zone or a local hospital
 - d. Consider ALS ground intercept be dispatched
 - i. Patient condition indicates shock
 - ii. Reports or confirmation of Penetrating and/or Significant Blunt Trauma to head, neck or torso
 - iii. Reports or confirmation of amputated extremities above fingers/toes, and/or multiple long bone fractures
 - iv. Reports or confirmation of burns
 - v. Reports or confirmation of toxic inhalation injury
 - vi. Reports or confirmation of entrapment with extrication/rescue needed
 - vii. Reports or confirmation of decreased or loss of consciousness
 - viii. Reports or confirmation of a multiple patient incident
 - ix. Injuries that may require pain management
 - x. ALS may arrive sooner by ground then by Air Ambulance
 - xi. Helicopter Air Ambulance unavailable or for other reason is not an option for this call
2. Notify Medical Control of Trauma Patient and possibility of trauma center candidate as early as possible

3. Consider Trauma System Activation and consult with medical control;
 - a. Vitals and LOC
 - i. Heart Rate >130
 - ii. Systolic BP <85
 - iii. Respiratory rate <10 or >29
 - iv. GCS <13
 - b. Anatomy of Injury
 - i. Penetrating Trauma to head, neck, torso, groin
 - ii. Combinations of burns >20% or face/airway burns
 - iii. Amputation above wrist/ankle
 - iv. Spinal Cord Injury
 - v. Flail Chest
 - vi. Two or more proximal long bone injuries
 - c. Biomechanics of injury
 - i. Ejected from Vehicle
 - ii. Auto vs Pedestrian/Bicycle >5 mph
 - iii. Motorcycle/ ATV crash
 - iv. Pedestrian thrown or run over
 - d. Other Risk Factors
 - i. Provider impression
 - ii. Extreme(s)
 1. age (<2 >60)
 2. environment (heat/cold)
 3. Health/Illness (Pregnancy, COPD, CHF, Diabetes)
 4. Haz/Mat
 - iii. High Energy Transfer
 1. Rollover
 2. Fall >10 feet
 3. Extractions > 20 minutes
 - iv. Burn Injury
 1. 2nd and 3rd degree burns of face, hands, feet, perineum
 2. significant electrical burns
 3. inhalation injury
4. Procedure
 - a. Consult with medical control
 - b. Advise patient condition and injuries
 - c. Request Trauma System Activation
 - d. Medical Direction approves trauma system activation
 - i. Transport patient to closest designated trauma center OR Local hospital for immediate stabilization.
 - e. If not already done consider ALS ground intercept or Helicopter Air Ambulance transport
5. Follow your Trauma Regions guidelines for transport of patient to the most appropriate facility and for more guidance on trauma system in your area.

The goal of the TRAUMA SYSTEM is to get the injured patient to the most appropriate facility by the most appropriate means in a timely manner. EMS needs to consult with Medical Control if any patient meets trauma system guidelines so the patient is transported to the most appropriate facility. In some cases the patient may bypass a local hospital or stop only to be stabilized by the local hospital then transferred on to a regional trauma center.

Head / Facial Injuries

First Responder

Scene Safety- BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen Consider assisting ventilations with BVM
Circulation	Vital Signs, Control External bleeding
Eye Trauma	Chemical exposure – Continuously Flush Eye Penetrating Object –Leave in Place, Stabilize with Dressings, Patch Other Eye Loss of Tissue-Keep Moist and Transport with Patient
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Airway	If No Gag Reflex, Insert Oral Airway DO NOT USE NASAL AIRWAY
Transport	Non-emergent Unless Patient Becomes Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Airway	If No Gag reflex, Insert Medical Director Approved Advanced Airway Device Multi-lumen Airway use with Caution if Facial Fractures Suspected
IV	Establish Peripheral IV Access

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Determine Cardiac Rhythm
Airway	Oral Intubation Only DO NOT NASAL INTUBATE

Paramedic

Assess	Perform Comprehensive Assessment
Monitor	O2 Sats and End Tidal CO2
Consider	Emergency Cricothyrotomy if Oral Intubation Cannot be Performed
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Brain Trauma Guidelines

Maintain

- a. O² Sat > 90%
- b. Systolic BP >90
- c. EtCO² 30 to 35mmHg

Soft Tissue Neck Injuries**First Responder**

Scene Safety- BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Control External Bleeding with Occlusive Dressing
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Airway	If No Gag Reflex, Insert Oral Airway If Gag Reflex May Use Nasal Airway if No Head Injury Suspected AND Patient has Decreased LOC
Transport	Non-emergent Unless Patient Becomes Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Airway	If No Gag reflex, Insert Medical Director Approved Advanced Airway Device (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
--------	--------------------------------------

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Emergency Cricothyrotomy if Oral Intubation Cannot be Performed
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Chest Injuries

First Responder

Scene Safety- BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen Seal Sucking Chest Wounds with Occlusive Dressing Remove Occlusive Dressing if Patient's Breathing Deteriorates
Circulation	Vital Signs,
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Airway	If No Gag reflex, Insert Oral Airway If Gag Reflex may use Nasal Airway if No Head Injury Suspected AND Patient has Decreased LOC
Transport	Non-emergent Unless Patient Becomes Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Airway	If No Gag reflex, Insert Medical Director Approved Advanced Airway Device (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Consider	Needle Decompression for Tension Pneumothorax

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Rapid Sequence Intubation (RSI) *See <u>RSI</u> Protocol

Abdominal /Pelvic Injuries**First Responder**

Scene Safety- BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs Control External Bleeding Seal Abdominal Eviscerations with Occlusive Dressing DO NOT REPLACE/REINSERT ABDOMINAL CONTENTS
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Consider	Splint Unstable Pelvis with PASG
Transport	Non-emergent Unless Patient Becomes Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
Consider	Cardiac Monitoring

Paramedic

Assess	Perform Comprehensive Assessment
--------	----------------------------------

Extremity Injuries

First Responder

Scene Safety- BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs Control External Bleeding
Assess	Conduct a Simple Patient Assessment
Splint	Splint Extremity Deformities Assess Pulse Distal to Injury Prior To and After Splint is Applied
Ice/elevate	Apply Ice Pack and If Possible Elevate the Extremity

EMT

Assess	Conduct Basic Patient Assessment
Monitor	Pulse distal to injury, <u>Suspected Fractures:</u> If Pulse Absent Follow Listed Steps, Stopping at the Step which Pulse Returns; 1 st Loosen Splint Recheck Pulse 2 nd Gently Reposition Limb/Straighten 3 rd Apply Gentle Traction to Limb 4 th Contact Medical Control Suspected Dislocations: If Pulse Absent Contact Medical Control
Consider	Splint Unstable Pelvis with PASG or Back Board with Sheet Wrap or Scoop Type Stretcher
Transport	Non-emergent Unless Patient Becomes Unstable OR Patient has No Pulse Distal to Injury
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
Consider	Pain Management *See <u>Pain Management</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Pain Management *See <u>Pain Management</u> Protocol

Ingested Poisons**First Responder**

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment – Gain Information About the Ingested Poison Product/Medication Name, Amount Ingested, Time of Ingestion
Contact	Poison Control Center According to Local Procedure

EMT

Assess	Conduct Basic Patient Assessment
Consider	Oral Activated Charcoal if Patient Can Maintain His/Her Own Airway
Transport	Non Emergent Unless Patient Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Consider	Inserting Medical Director Approved Advanced Airway Device (Multi-lumen Airway, LMA, ET)
Consider	Obtain Glucose Reading, And if Indicated Oral Glucose <u>AND</u> if Patient's Airway can be Maintained
IV	Establish Peripheral IV Access

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Suspected	If Poison is Identified or a High Suspicion *See <u>Specific Agent/Toxin</u> Protocols
Consider	Dextrose 50% 25g IVP If Indicated by Glucometer Reading
Consider	Narcan 2mg IV If Narcotic Drug Intake is Suspected
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol

Specific Agents/Toxins
Known or High Suspicion of Narcotic Overdose
This protocol is for the EMT-I 99 and Paramedic

Basic Life Support

BLS Measures as Outlined in the Ingested Poisons Protocol

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Airway	If Airway can be Managed with Manual Maneuvers and Suction delay Advanced Airway until Narcan is Given.
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Consider	Naloxone .4mg to 2mg IV
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol
Assess	If Patient Condition does not Improve Consider Other Possible Causes
Prepare	The Patient May Have to be Restrained Once the Effects of the Narcotic is Reversed

Paramedic

Assess	Perform Comprehensive Assessment
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol

***Narcotic overdoses may require additional doses of naloxone to maintain the patient's vital signs and LOC.
The patient must be closely monitored.

Specific Agents/Toxins
Known or High Suspicion of Tricyclic Anti-Depressant Overdose
This protocol is for the EMT-I 99 and Paramedic

Basic Life Support

BLS Measures as Outlined in the Ingested Poisons Protocol

EMT-Intermediate 99

- | | |
|--------------------|---|
| Assess | Perform Advanced Physical Assessment |
| Consider | Inserting Medical Director Approved Advanced Airway Device
(Multi-lumen Airway, LMA, ET) |
| Cardiac
Monitor | Attach Cardiac Monitor, Interpret ECG |
| Dysrhythmia | *See <u>Cardiac Dysrhythmia</u> Protocol |
| Consider | 500 cc Fluid Challenge
For Hypotension |

Paramedic

- | | |
|-------------|--|
| Assess | Perform Comprehensive Assessment |
| Dysrhythmia | *See <u>Cardiac Dysrhythmia</u> Protocol |
| Consider | Sodium Bicarbonate
1mEq/kg, slow IVP |

Specific Agents/Toxins
Known or High Suspicion of Organophosphate OR Nerves Agent Exposure
This protocol is for the EMT-I 99 and Paramedic

Basic Life Support

The patient must be removed from the hazard area and decontaminated by trained rescuers
BLS measures as outlined in the Ingested Poisons Protocol

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Inserting Medical Director Approved Advanced Airway Device (Multi-lumen Airway, LMA, ET)
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol
Consider	Atropine 2mg IV Every 5 Minutes Until Symptoms Relieved
Consider	Diazepam 5-10mg for Seizures

Paramedic

Assess	Perform Comprehensive Assessment
Dysrhythmia	*See <u>Cardiac Dysrhythmia</u> Protocol
Alternate Medication	Lorazepam 2-4mg May be Given for Seizures as Alternate to Diazepam

Specific Agents/Toxins
Known Calcium Channel Blocker Overdose
This protocol is for the EMT-I 99 and Paramedic

Basic Life Support

BLS measures as outlined in the Ingested Poisons Protocol

EMT-Intermediate 99

Assess

Perform Advanced Physical Assessment

Consider

Inserting Medical Director Approved Advanced Airway Device
(Multi-lumen Airway, LMA, ET)

Cardiac
Monitor

Attach Cardiac Monitor, Interpret ECG

Dysrhythmia

*See Cardiac Dysrhythmia Protocol

Paramedic

Assess

Perform Comprehensive Assessment

Consider

Calcium Chloride 500 – 1000mg Slow IV

Dysrhythmia

*See Cardiac Dysrhythmia Protocol

Toxic Inhalation**First Responder**

Scene Safety –
BSI

SAFETY FIRST
PATIENT SHOULD BE REMOVED FROM HAZARD AREA BY
TRAINED RESCUERS AND IF REQUIRED DECONTAMINATED

Level of
Conscious

Alert, Verbal, Pain, or Unresponsive

Airway

Monitor Airway

Breathing

Administer Oxygen

Consider

Assisting Ventilations with BVM

Circulation

Vital Signs, Skin Color/Temp

Assess

Conduct a Simple Patient Assessment

EMT

Assess

Conduct Basic Patient Assessment

Consider

Assisting Patient with His/Her Metered Dose Inhaler

Transport

Non –Emergent Transport Unless Patient is Unstable

Consider

ALS Intercept

EMT Options/ EMT Intermediate 85

IV

Establish Peripheral IV Access

EMT-Intermediate 99

Assess

Perform Advanced Physical Assessment

Consider

Bronchodilator Medication by Nebulizer
*See Bronchodilator Protocol

Cardiac
Monitor

Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess

Perform Comprehensive Assessment

Consider

Rapid Sequence Intubation (RSI)
See RSI Protocol

Burns

First Responder

Scene Safety- BSI	SAFETY FIRST
Stop	Burning Process, Remove Heat Source if Possible Cool Burning Material Adhering to the Patient
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs
Assess	Conduct a Simple Patient Assessment
Prevent	Hypothermia
Remove	Jewelry or Other Restrictive Items From or Near Burn Area Burned Clothing, Cut Around Clothing Adhering to the Skin
Estimate	Body Surface Area (BSA) Burned
Dress Burns	Cover Burn Area with Dry Bandages or Sheets If BSA <10% May Cool Burn Area PREVENT HYPOTHERMIA

EMT

Assess	Conduct Basic Patient Assessment
Transport	Emergent for Burns to Face/Airway Emergent for >30 Minutes Transport Time with No ALS Burns >10%BSA
Consider	ALS Intercept or ALS Helicopter transport for any burns to face/airway

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
Consider	Pain Management *See <u>Pain Management</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Pain Management *See <u>Pain Management</u> Protocol
Consider	Rapid Sequence Intubation (RSI) for Facial/Airway burns ***See <u>RSI</u> Protocol

Snake Bite

First Responder

Scene Safety- BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs
Calm	Calm Patient, Keep Patient still Keep Extremity BELOW the Level of the Heart
Rinse	The Bite Site DO NOT RUB or SCRUB DO NOT APPLY ICE, DO NOT CUT/INCISE BITE
Apply	Restrictive Band 1 Inch above the Bite. The Pulse Should be Palpable Distal to the Band
Assess	Conduct a Simple Patient Assessment
Identify	If the Snake is Dead, Transport it to the Hospital **Warning - The Bite Reflex is Active Even After the Snake is Dead If The Snake is Not Present Investigate to Find Out the Type of Snake

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non-emergent Unless Patient Becomes Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT- Intermediates 99

Assess	Perform Advanced Physical Assessment
Consider	Pain Management *See <u>Pain Management</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Pain Management *See <u>Pain Management</u> Protocol

PEDIATRICS

Cardiopulmonary Arrest

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Confirm unresponsiveness
AED	Attach AED If Patient Older then 8 and Over 60 lbs ** SEE AED Protocol***
Airway	Establish an Airway
Breathing	Ventilate with Bag Valve Mask Attached to O2 OR Pocket Mask
Circulation	Begin Chest Compressions if Pulseless OR Bradycardia (< 80 beats/min in newborn or < 60 beats/min in infants) AND Do Not Respond to Ventilation and Oxygenation
Prepare	Package for transport

EMT

Airway	Insert an Oral Airway
Assess	Patient's Weight
Transport	Emergent
Consider	ALS Intercept

EMT Options / EMT Intermediate 85

Airway	Insert Medical Director Approved Advanced Airway Device (Multi-lumen Airway, LMA, ET) Multi-lumen Airways Only for Patients 5'2" or Taller
IV	Establish Peripheral IV Access

EMT Intermediate 99

Consider	IO Access In Lieu of IV Access
PALS	Follow Appropriate <u>Pediatric Cardiac Arrest</u> Algorithm

Paramedic

PALS	Follow Appropriate <u>Pediatric Cardiac Arrest</u> Algorithm
------	--

Note: If the patient regains a pulse see the Pediatric Post Cardiac Arrest – Return of Pulse protocol

Pediatric Cardiac Arrest Algorithm**V-Fib****EMT-Intermediate 99**

Shock	3 Shocks 2J/Kg, 4J/Kg, 4J/Kg
Airway	Establish an Airway Endotracheal Intubation Preferred
Breathing	Ventilation with Bag Valve Mask attached to O2
Circulation	Chest Compression and Establish Peripheral IV Access
Medication	Epinephrine .01mg/Kg (1:10,000) IV/IO Every 3-5 Minutes Or Epinephrine .1mg/Kg (1:1,000) ET
Shock	Shock 4J/Kg
Intervention Sequence	Drug Shock 4J/Kg Repeat
Medication Considerations	Lidocaine 1 mg/kg

If IV access is delayed or can not be obtained Epi may be given via the ET tube

If the patient's cardiac rhythm changes see appropriate protocol for that rhythm.
If patient develops a pulse see the Pediatric Post Cardiac Arrest protocol.

Paramedic

Medication Considerations	Amiodarone 5mg/Kg
	Magnesium Sulfate 25-50mg/kg to Max of 2g For torsades de pointes

Pediatric Cardiac Arrest Algorithm PEA

EMT-Intermediate 99

Confirm	Cardiac Rhythm
Airway	Establish Airway Endotracheal Intubation Preferred
Breathing	Ventilation with Bag Valve Mask attached to O2
Circulation	Chest Compression and Establish Peripheral IV Access
Medication	Epinephrine .01mg/Kg (1:10,000) IV/IO Every 3-5 Minutes Or Epinephrine .1mg/Kg (1:1,000) ET
Intervention Sequence	Drug Evaluate for change Repeat
Medication Considerations	Epinephrine .1mg/Kg (1:1,000) ET, IV, IO Every 3-5 Minutes
Consider	Consider Causes Hypovolemia Consider Fluid Boluses 20cc/Kg Tension Pneumothorax Consider Needle Decompression Hypothermia Consider Warming Patient

If IV access is delayed or can not be obtained Epi may be given via the ET tube

If the patient's cardiac rhythm changes see appropriate protocol for that rhythm. If patient develops a pulse see the Pediatric Post Cardiac Arrest protocol.

Paramedic

Consider	Consider causes Acidosis Consider Sodium Bicarbonate Tricyclic Overdose Consider Sodium Bicarbonate Calcium Channel Blocker Overdose Consider Calcium Chloride
Medication Doses	Sodium Bicarbonate 8.4% 1mEq/Kg (4.2% in Neonates) Calcium Chloride 20 to 25 mg/Kg

Pediatric Cardiac Arrest Algorithm

Asystole

EMT-Intermediate 99

Confirm	Cardiac Rhythm
Airway	Establish Airway Endotracheal Intubation Preferred
Breathing	Ventilation with Bag Valve Mask Attached to O2
Circulation	Chest Compression and Establish Peripheral IV Access
Medication	Epinephrine .01mg/Kg (1:10,000) IV/IO every 3-5 Minutes Or Epinephrine .1mg/Kg (1:1,000) ET
Intervention Sequence	Drug Evaluate for change Repeat

If IV access is delayed or can not be obtained Epi may given via the ET tube

Medication Considerations	Epinephrine .1mg/Kg (1:1,000) ET, IV, IO Every 3-5 minutes
------------------------------	---

Consider	Consider Causes Hyoethermia Consider Warming Patient
----------	---

Paramedic

Consider	Consider Causes Acidosis Consider Sodium Bicarbonate Tricyclic Overdose Consider Sodium Bicarbonate Calcium Channel Blocker Overdose Consider Calcium Chloride
Medication Doses	Sodium Bicarbonate 8.4% 1mEq/Kg (4.2% in Neonates) Calcium Chloride 20 to 25 mg/Kg

If the patient's cardiac rhythm changes see appropriate protocol for that rhythm.
If patient develops a pulse see the Pediatric Post Cardiac Arrest protocol.

Pediatric Post Cardiac Arrest – Return of Pulse

With access to AEDs the OOH care provider may respond to a person who has been successfully defibrillated OR the OOH provider may successfully defibrillate/resuscitate a cardiac arrest victim.

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Confirm Unresponsiveness
Airway	Establish an Airway
Breathing	Assist Ventilations with Bag Valve Mask
Circulation	Confirm Pulse Present, Recheck Often
Assess	Conduct Simple Patient Assessment

EMT

Airway	Insert Oral Airway
Assess	Perform Basic Patient Assessment
Transport	Emergent
Consider	ALS Intercept

EMT Options / EMT Intermediate 85

Airway	Insert Medical Director Approved Advanced Airway Device Appropriate for Age and Size (Multi-lumen Airway, LMA, ET)
IV	Establish Peripheral IV Access
Consider	20cc/kg Fluid Bolus if Hypotensive for Age

EMT Intermediate 99

Consider	IO access In Lieu of IV Access
Assess	Perform Advanced Assessment
Cardiac Monitor	Determine Cardiac Rhythm
Dysrhythmias	Treat with Appropriate <u>Pediatric Advance Cardiac Dysrhythmia</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
Dysrhythmias	Treat with Appropriate <u>Pediatric Advanced Cardiac Dysrhythmia</u> Protocol

Pediatric General Cardiac Dysrhythmia

The First Responder, EMT, EMT with Option, and EMT-I 85 will only detect by evaluation of the pulse whether the pulse is fast. Slow. Or normal rate and if it is regular or irregular. EMT-I 99 and Paramedic will interpret the cardiac rhythm. Generally pediatric patients do not have cardiac dysrhythmias due to cardiac disease, most often the cause is an airway/ventilation problem or volume problem. The OOH provider should secure an airway and ventilate with a BVM for the pediatric patient who shows the signs and symptoms of poor perfusion.

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Painful, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Consider	Assisting Ventilations with BVM
Circulation	Assess Pulse Rate, Rhythm, and Quality Vital Signs <u>Begin chest compressions if heart rate <60/min in infants</u>
Assess	Conduct Simple Patient Assessment
Prepare	Prepare patient for transport

EMT

Assess	Perform a Basic Assessment
Transport	Non-emergent Transport Unless Patient Unstable
Consider	ALS Intercept

EMT Options / EMT Intermediate 85

Airway	Insert Medical Director Approved Advanced Airway Device (Multi-lumen Airway, LMA, ET) Multi-lumen airways only for patient 5'2" or taller
IV	Establish Peripheral IV Access

EMT Intermediate 99

Consider	IO access in lieu of IV access
Assess	Perform Advanced Assessment
Cardiac Monitor	Determine Cardiac Rhythm
PALS	Follow Appropriate <u>Pediatric Advanced Cardiac Dysrhythmia</u> Protocol

Paramedic

Assess	Perform Comprehensive Assessment
PALS	Follow Appropriate <u>Pediatric Advanced Cardiac Dysrhythmia</u> Protocol

Pediatric Advanced Cardiac Dysrhythmia

This Protocol applies to EMT-I 99 and Paramedics and assumes the General Cardiac Dysrhythmia protocol has been followed

For the stable patient tolerating the cardiac rhythm may require monitoring and supportive care without a medication or electrical intervention.

For the unstable patient the EMT-I 99 or Paramedic may have to consider a more aggressive treatment plan based on the patient condition and how rapidly a medication may be delivered verses an electric therapy can be performed.

Bradycardia with signs and symptoms of poor perfusion

EMT-Intermediate 99

Assess/ Reassess	Reassess Airway and Ventilations Secure Airway and Assist Ventilations
Consider	Epinephrine .01mg/Kg IV(1:10,000) or .1mg/kg ET (1:1000)
Consider	Atropine .02mg/Kg Minimum Dose .1mg Max Dose .5mg Child or 1mg Adolescent
Consider	Medication not effective Transcutaneous Pacing Premedicate if Possible Diazepam .25mg/Kg

Paramedic

Alternate Pre-medications	Midazolam 0.1mg/kg iv to max of 2.5mg Lorazepam .05 to .2/mg/Kg
Consider	Epinephrine Infusion .1 -1 mcg/Kg/Min OR Dopamine 2 – 20 mcg/Kg/Min

Ventricular Tachycardia

EMT Intermediate 99

Assess/ Reassess	Reassess Airway and Ventilations Secure airway and assist ventilations
Consider	12lead ECG
Consider	Synchronized Cardioversion .5-1J/Kg Premedicate if Possible Diazepam .25mg/Kg
If Dysrhythmia Resolves	Lidocaine 1mg/kg bolus followed by Lidocaine Infusion 20-50mcg/Kg/Min/min

Paramedic

Alternate Pre-medications	Midazolam 0.1mg/kg iv to max of 2.5mg
------------------------------	---------------------------------------

**Pediatric Advanced Cardiac Dysrhythmia
Continued****PSVT with sign and symptoms of poor perfusion****EMT Intermediate 99**

Consider

Synchronized Cardioversion
Premedicate if Possible
Diazepam .25mg/Kg

OR

Consider

Adenosine 0.1-0.2 mg/Kg
Rapid IV push**Paramedic**Alternate
Pre-medications

Midazolam 0.1mg/Kg IV to Max of 2.5mg

Pediatric Difficulty Breathing
Asthma/Bronchiolitis
Difficulty Breathing in the presence of wheezing

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen Consider Assisting Ventilations with BVM
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

First Responder and EMT

Consider	Epinephrine Auto Injector Pediatric (EPI PEN Jr) for Impending Respiratory Collapse Guidelines; Patient Able to speak one-two word phrases Low/falling oxygen saturations even with O2 administration Diminished to absent lung sounds Decreasing LOC Retractions Pale or cyanotic skin
----------	---

EMT

Assess	Conduct Basic Patient Assessment
Consider	Assist Patient with His/Her Metered Dose Inhaler (MDI)
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access
----	--------------------------------

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Bronchodilator Medication by nebulizer *See <u>Bronchodilator</u> Protocol
Consider	IO Access In Lieu of IV Access Do Not Delay Epi to Obtain IV/IO Access
Consider	Epinephrine (1:1,000) 0.01 mg/kg SQ, Max Dose 0.3 mg. May Repeat Every 5 to 10 Minutes
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Rapid Sequence Intubation (RSI) ***See <u>RSI</u> Protocol***

Laryngotracheobronchitis (Croup), Epiglottitis

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Approach	In Calm Manner DO NOT Excite/Scare the Child
Airway	Monitor Airway
Breathing	Administer Oxygen EXTREME CAUTION MUST BE EXERCISED Epiglottitis may cause the patient airway to become occluded completely if the patient is agitated
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

IV	Establish Peripheral IV Access EXTREME CAUTION MUST BE EXERCISED Epiglottitis may cause the patient airway to become occluded completely if the patient is agitated in the process of IV insertion
----	--

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Bronchodilator Medication by Nebulizer if Laryngotracheobronchitis (Croup) is Suspected *See <u>Bronchodilator</u> Protocol
Consider	IO Access In Lieu of IV Access
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess	Perform Comprehensive Assessment
Consider	Cricothyrotomy Needle or Surgical for Occluded Airway

Acute Allergic Reaction / Anaphylaxis**Difficulty Breathing in the presence of urticaria, wheezing and /or contact with a known allergen****First Responder**

Scene Safety –

BSI SAFETY FIRST

Level of

Conscious Alert, Verbal, Pain, or Unresponsive

Airway

Monitor Airway

Breathing

Administer Oxygen,

Consider

Assisting Ventilations with BVM

Circulation

Vital Signs, Skin Color/Temp

Assess

Conduct a Simple Patient Assessment

Consider

Epinephrine Auto Injector Pediatric (EPI PEN Jr) if Impending Respiratory Collapse Guidelines; Patient able to speak only one-two word phrases without taking a breath
 Low/falling oxygen saturations even with O2 administration
 Diminished to absent lung sounds
 Decreasing LOC
 Retractions
 Pale or cyanotic skin

EMT

Assess

Conduct Basic Patient Assessment

Consider

Assist Patient with His/Her Metered Dose Inhaler

Transport

Emergent Transport Unless Patient is stable

Consider

ALS Intercept

EMT Options/ EMT Intermediate 85

IV

Establish Peripheral IV Access Titrate to Blood Pressure

EMT-Intermediate 99

Consider

Bronchodilator Medication by Nebulizer *See Bronchodilator Protocol

Consider

IO Access In Lieu of IV Access
 Do Not Delay Epi for IV/IO Access

Consider

Epinephrine (1:1,000) 0.01 mg/kg SQ for bronchospasm, maximum dose 0.3 mg.
 May Repeat Every 5 Minutes if Severe Respiratory Symptoms are Not Resolved

Cardiac

Monitor

Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess

Perform Comprehensive Assessment

Consider

Consider Diphenhydramine (Benadryl), 1 mg/kg, IM or slow IVP over 1-3 min.
 Maximum Individual Dose 50mg.

Consider

Rapid Sequence Intubation (RSI) ***See RSI Protocol***

Pediatric Upper Airway Obstruction

First Responder

Scene Safety –
BSI

SAFETY FIRST

Level of
Conscious

Alert, Verbal, Pain, or Unresponsive

Airway

Attempt to Relieve Obstruction Using AHA Guideline for Obstructed Airway for Pediatric Patients

Continue

Obstruction not Cleared
Attempts to Relieve
Obstruction

Obstruction Cleared

Breathing

Administer Oxygen
Consider Assisting Ventilations
with BVM

Circulation

Vital Signs, Skin Color/Temp

Assess

Conduct a Simple Patient Assessment

EMT

Transport

Emergent
Continue Attempts to Relieve
Obstruction

Non-emergent Transport if Patient Stable

Assess

Conduct Basic Patient Assessment

Consider

ALS Intercept

ALS Intercept

EMT Options/ EMT Intermediate 85

Airway

Attempt to Visualize Obstruction with
Laryngoscope and Remove with McGill
Forceps. IF TRAINED

Consider Advance Airway if Patient
LOC Remains Decreased and No Gag Reflex

IV

Establish Peripheral IV Access
Do Not Delay Transport

Establish Peripheral IV Access

EMT-Intermediate 99

Consider

IO Access In Lieu of IV Access

Assess

Perform Advanced Physical Assessment

Cardiac Monitor

Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess

Perform Comprehensive Assessment

Consider

Cricothyrotomy needle or surgical

Pediatric Seizures

First Responder

Scene Safety – BSI	SAFETY FIRST
Level of Conscious	Alert, Verbal, Pain, or Unresponsive
Airway	Monitor Airway
Breathing	Administer Oxygen
Circulation	Vital Signs, Skin Color/Temp
Assess	Conduct a Simple Patient Assessment
Prepare	Package Patient for Transport

EMT

Assess	Conduct Basic Patient Assessment
Transport	Non –Emergent Transport Unless Seizures Continue
Consider	ALS Intercept

EMT Options/ EMT Intermediate 85

Glucometer	Obtain Glucose Reading
IV	Establish Peripheral IV Access

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Consider	Possible Causes of Seizure
Consider	Diazepam .25mg/Kg IV to Max of 5mg for Recurrent or Prolonged Seizures
Consider	For Age 1 and Less Dextrose 25% .5 to 1gm/Kg if Glucometer Reading Indicated For Age over 1 Dextrose 50% .5 to 1 gm/Kg if Glucometer Reading Indicated
Cardiac Monitor	Attach Cardiac Monitor, Interpret ECG

Paramedic

Assess	Perform Comprehensive Assessment
Alternate Medication	Lorazepam .05 -.2 mg/Kg IV, IO

Appendixes
Bronchodilator Protocol

EMT, EMT Options, and EMT Intermediate 85

Assist May Assist Patient with His/Her Metered Dose Inhaler (MDI)
 Should be administered as prescribed
 *Note : Not All MDI's are for emergent situations for patients suffering
 acute shortness of breath
 Listed below are common emergent medications

EMT-Intermediate 99 and Paramedic

Consider Bronchodilator medication by nebulizer

Medication Options:

Generic Name	Trade Name	Nebulizer dose
Albuterol	Proventil, Ventolin	2.5mg in 3cc NS
Ipratropium	Atrovent	.5mg (500mcg) in 2 to 3 cc NS
Albuteral/Inpatropium	Combivent, Duoneb	3mg Albuterol .5mg Inpatropium in 3cc NS
Metaproterenol	Alupent	.2-.3 cc of 5% solution

Pain Management

1st Responder, EMT

Position	Unless Otherwise Contraindicated Because of Trauma Place Patient in Position of Comfort
Splint	Apply Splint to Extremity Deformities Apply Ice and Elevate to Reduce Pain

EMT Options/ EMT Intermediate 85

IV	Establish IV Access
----	---------------------

EMT-Intermediate 99

Consider	Morphine 2-5 mg IV or IM [Pediatric Dose .1-.2 mg/kg] May Repeat in 2mg Doses Until Pain Control or 10mg Total and BP Remains >100 Systolic
----------	--

Paramedic

Consider	The Listed Medications for Pain Control/Management
----------	--

Medication Option	Adult Dose	Pediatric Dose
Morphine	2-5mg	.1 -.2mg/Kg
Fentanyl	25 – 100 mcg	1 –4 mcg/Kg
Meperidine *	50 – 100 mg	1mg/Kg
Ketorolac	30mg IV 60mg IM	.5 – 1 mg/Kg to 30mg max IV

* Note: When considering Meperidine also consider the use of an anti-emetic such as Phenergan

* Note: For advanced providers the listed medications with the exception of Ketorlac are control substances and all medication administrations and wastes of unused medications must be documented per local policy.

Hospital to Hospital Transfer Protocol

EMT

Scene Safety – BSI	SAFETY FIRST
Assess	Conduct Basic Patient Assessment
Oxygen	Continue oxygen therapy with Nasal Cannula or Non-Rebreather Mask
Report	Contact Patient's Nurse for a Patient Report
Transport	Non –Emergent Transport Unless Patient is Unstable
Consider	ALS Intercept if Patient Becomes Unstable

EMT Options/ EMT Intermediate 85

Airway	If Advanced Airway Placed Confirm Correct Placement
IV	Monitor Established IV of an Isotonic Solution at the Transporting Facilities Ordered Rate

EMT-Intermediate 99

Assess	Perform Advanced Physical Assessment
Cardiac Monitor	Cardiac Monitoring for All Cardiac Patients and All Unstable Patients
Medications	The EMT-I 99 May Only Administer the Medications As listed In the Rules and Regulations Practices and Procedures for the EMT Intermediate. (Title 172 NAC 11)
Infusions	The EMT-I 99 May Only Establish/Maintain an Infusion of Lidocaine No Other Infusions of Medications are Allowed by this Protocol.

Paramedic

Assess	Perform Comprehensive Assessment
Medications	The Paramedic may with a Physician Order From Either the Transferring or Receiving Facility Administer Medication(s) by Any Ordered Route.
Infusions	The Paramedic may with a Physician Order from Either the Transferring or Receiving Facility Maintain, Adjust, or Initiate an Ordered Infusion.

***Follow appropriate protocol for a specific problem if the patient condition changes or new sign/symptoms are recognized by the provider. On line medical control should be consulted and advised if patient becomes unstable.

Anti Emetic Protocol
For Paramedic level providers only

Guidelines for Anti Emetic treatment:

- Nausea/Vomiting from Trauma/Pain
- Nausea/ Vomiting due to Medication Side Effect
- Nausea/Vomiting from an Acute Cardiac or Medical Event
- Nausea/Vomiting due to Ambulance Transport

Assess and Treat the Cause of Nausea if Possible

Follow Appropriate Protocol for the Patient's Condition

Administer One of the Medications Listed as Approved by the PMD

Medication choices

Medication	Dose
Anzemet	12.5mg IV
Compazine (prochlorperazine)	5-10mg IV
Phenergan (promethazine)	12.5-25 IV
Zofran (ondansetron)	4-8mg IV

Rapid Sequence Intubation

For Paramedic level providers only

Criteria for Rapid Sequence Intubation

GCS of ≤ 8

Respiratory failure/impending failure

Patient unable to protect his/her own airway

Head injuries with decreased LOC/ combativeness s/s of increased ICP

Consideration before attempting RSI:

Benefit vs Risk of procedure

Difficulty of intubation

Backup Airway Plan if Intubation Fails

Steps	Details
Preoxygenate	1. Preoxygenate with 100% oxygen by mask. If ventilatory assistance is necessary, ventilate gently, applying cricoid pressure.
Premedicate	2. Premedicate as appropriate; then WAIT 3 MINUTES after drug administration. <ul style="list-style-type: none"> • Sedation with Diazepam: 5 to 10 mg (pediatric dose 0.25mg/kg not to exceed 5mg) or Versed 0.1 to 0.2 mg/kg titrated (max dose 2mg) • Atropine: 0.01 mg/kg IV push for children or adolescents (minimum dose of 0.1 mg recommend) • Lidocaine: 1.0 to 1.5 mg/kg IV over 30 to 60 seconds • Defasciculating agent (optional, see Table 2)
Sedate then Paralyze	3. Induce sedation with one of these agents: diazepam, thiopental, fentanyl, ketamine, etomidate, versed, or methohexital. (See Table 1) 4. Give succinylcholine 1.5 mg/kg IV push (use 2.0 mg/kg for infants and small children). 5. Assess for apnea, jaw relaxation, decreased resistance to bag-mask ventilations (patient sufficient relaxed to proceed with intubation). 6. Apply cricoid pressure.
Placement: performance	7. Perform tracheal intubation. If unable to intubate within 20 seconds, stop. Ventilate with bag-mask for 30 to 60 seconds. Use pulse oximetry as a guide. Inflate balloon cuff when TT is in place.
Placement: primary confirmation	9. Perform primary confirmation of TT placement: <ul style="list-style-type: none"> • By direct visualization of TT passing through vocal cords • By chest rise/fall with each ventilation (bilateral) • By 5-point auscultation: anterior chest L and R, midaxillary line L and R, and over the epigastrium (Listen for air entering the stomach when BM is squeezed and by tube condensation.)
Placement: secondary confirmation	10. Perform secondary confirmation of TT placement: <ul style="list-style-type: none"> • Use a bulb aspiration device(esophageal detector device [EDD]) • If the EDD indicates that the TT is in the trachea, leave in place monitor O2 sat • Monitor end-tidal CO2
Placement: prevent dislodgement	11. Secure TT with commercial holder (preferred) <ul style="list-style-type: none"> • Alternatively, secure with an adhesive tape/cloth cord technique. • In out-of-hospital setting with the prospect of patient ventilation during movement, immobilize cervical spine with cervical collar or backboard or both.
Maintain Sedation/ Paralytic	12. Administer maintenance dose of paralytic agent 13. Administer sedative if transport time longer then duration of sedative agent 14. Consider pain management agent (Morphine, or Fentanyl)

Rapid Sequence Intubation

Table 1
Sedative and Induction Agents

Sedative	Dosage IV Push	Onset	Duration
Etomidate	0.2 to 0.6 mg/kg	60 seconds	3 to 5 minutes
Fentanyl	Induction: 2 to 10 ug/kg Sedation (titrate): 2 to 4 ug/kg	60 seconds	30 to 60 minutes
Ketamine	2.0 mg/kg	30 to 60 seconds	15 minutes
Midazolam (Versed)	Induction: 0.07 to 0.3 mg/kg Sedation (titrate): 0.02 to 0.04 mg/kg	2 minutes	1 to 2 hours
Thiopental	3 to 5 mg/kg	20 to 40 seconds	5 to 10 minutes
Diazepam	5-10 mg	60-90 seconds	60 to 180 minutes
Methohexital (Brevital)	1-1.5 mg/kg	60 sec	5 to 7 minutes

Table 2
Neuromuscular Blocking Agents

Agent	Dosage (Paralytic)	Dosage (defasciculating)	Onset	Duration
Succinylcholine	RSI: 1 to 2 mg/kg		30 to 60 seconds	4 to 6 minutes
Vecuronium	RSI: .1 mg/kg M: 0.01-.05 mg/kg	0.01 mg/kg	2.5 to 5 minutes	25 to 40 minutes

RSI = Rapid Sequence Intubation

M = Maintenance dose